

Antibiotic Resistance

- Primary Care: facts and figures



Fact #1

Antibiotic resistance is an increasingly serious patient safety and public health problem in England, Europe and the world [1, 2]. Resistance to antibiotics is increasing. In many countries, resistance rates have more than doubled in five years (Figure 1).

Growing antibiotic resistance threatens the effectiveness of antibiotics now and in the future. Inappropriate use of antibiotics not only leads to resistance which increases the risk of treatment failure, hospitalisation and cost of care but can also lead to adverse effects such as thrush, and diarrhoea caused by *Clostridium difficile* infection




Figure 1. Trends in antibiotic resistance (invasive infections), 2002-2008. Source: EARSS, 2009 [3].

Fact #2

Antibiotic exposure leads to emergence of antibiotic resistance [4]. The overall uptake of antibiotics in an individual and the population, as well as how antibiotics are consumed, has an impact on antibiotic resistance [5, 6, 7]. The presence of antibiotic resistant bacteria for urinary, respiratory and skin infections in individual patients has been strongly linked to the number and duration of antibiotic courses prescribed in the previous 12 months [7].





Figure 2. The link between antibiotic consumption and antibiotic resistance, 2007 (or latest available year, each data point represents one country) [3,11]

Antibiotic Resistance

– Primary Care: facts and figures

Fact #3

In England, primary care accounts for about 80 % of all antibiotic prescriptions, mainly for respiratory tract infections [5]. Antibiotic prescribing and consumption varies between GP practices in England and between European countries [8-11]. Although there has been a reduction in prescribing of fluoroquinolones and cephalosporins, total usage of antibiotics in England continues to rise (Figure 3).




Figure 3. Trends in usage of antimicrobials in general practice in England (Courtesy NHSBSA)

http://www.nhsbsa.nhs.uk/PrescriptionServices/Documents/PPDPrescribingAnalysisCharts/Antibiotics_Jun_11

References

- [1] European Antimicrobial Resistance Surveillance System. EARSS Annual Report 2007. Bilthoven, Netherlands: National Institute for Public Health and the Environment, 2008. http://www.rivm.nl/earss/result/Monitoring_reports/.
- [2] Cars O, et al. Meeting the challenge of antibiotic resistance. *BMJ* 2008;337:a1438. doi: 10.1136/bmj.a1438.
- [3] European Antimicrobial Resistance Surveillance System (EARSS). Interactive database. <http://www.rivm.nl/earss/database/>.
- [4] Malhotra-Kumar S, et al. Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a randomised, double-blind, placebo-controlled study. *Lancet* 2007;369:482-90.
- [5] Standing Medical Advisory Committee Sub-Group on Antimicrobial Resistance 1998. The Path of Least Resistance. www.advisorybodies.doh.gov.uk/smac1.htm (Last accessed 08 August 2011).
- [6] Guillemot D, et al. Low dosage and long treatment duration of beta-lactam: risk factors for carriage of penicillin-resistant *Streptococcus pneumoniae*. *JAMA* 1998;279:365-70.
- [7] Costello C et al. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systemic review and meta-analysis. *BMJ* 2010; 340:c2096.
- [8] Mölstad S, et al. Antibiotic prescription rates vary markedly between 13 European countries. *Scand J Infect Dis* 2002;34: 366-71.
- [9] Goossens H, et al; ESAC Project Group. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet* 2005;365:579-87.
- [10] NHS Business Services Authority. National Antibiotics Charts - Prescription Services Information Services 2011. http://www.nhsbsa.nhs.uk/PrescriptionServices/Documents/PPDPrescribingAnalysisCharts/Antibiotics_Jun_11_-_National.pdf (08 August 2011, date last accessed).
- [11] European Surveillance of Antibiotic Consumption (ESAC). http://app.esac.ua.ac.be/public/index.php/en_gb.
- [12] Butler CC, et al. Understanding the culture of prescribing: qualitative study of general practitioners' and patients' perceptions of antibiotics for sore throats. *BMJ* 1998;317(7159):637-42.
- [13] Kallestrup P and Bro F. Parents' beliefs and expectations when presenting with a febrile child at an out-of-hours general practice clinic. *Br J Gen Pract* 2003;53(486):43-4.
- [14] Little PS, et al. An open randomised trial of prescribing strategies for sore throat. *B M J* 1997; 314:722-727.
- [15] Little PS et al. Re attendance and complications in a randomised trial of prescribing strategies for sore throat: the medicalising effect of prescribing antibiotics. *B M J* 1997; 315:350-352.
- [16] Moore M et al. Effect of antibiotic prescribing strategies and an information leaflet on longer-term reconsultation for acute lower respiratory tract infection. *B J G P* 2009; 567:728-734.
- [17] Macfarlane J, et al. Influence of patients' expectations on antibiotic management of acute lower respiratory tract illness in general practice: questionnaire study. *BMJ* 1997;315(7117):1211-4.
- [18] Li J, et al. Antimicrobial prescribing for upper respiratory infections and its effect on return visits. *Fam Med* 2009;41(3):182-7.
- [19] Rutten G, et al. Patient education about cough: effect on the consulting behaviour of general practice patients. *Br J Gen Pract* 1991; 41(348):289-92.
- [20] Cals JW, et al. Evidence based management of acute bronchitis: sustained competence of enhanced communication skills acquisition in general practice. *Patient Educ Couns* 2007;68(3):270-8.
- [21] Welschen I, et al. Effectiveness of a multiple intervention to reduce antibiotic prescribing for respiratory tract symptoms in primary care: randomised controlled trial. *BMJ* 2004; 329(7463):431-3

Fact #4

Communicating with patients is key.

Professional medical advice impacts patients' perceptions and attitudes towards their illness and perceived need for antibiotics, particularly if they are advised on what to expect in the course of the illness, including realistic recovery time and self-management strategies [12-18].

Patient satisfaction in primary care settings depends more on effective communication than on receiving an antibiotic prescription [12-18]. Prescribing an antibiotic for acute respiratory tract infections, medicalises illness and increases reconsultation rates [14-18].

Primary care prescribers do not need to allocate more time for consultations that involve offering alternatives such as delayed prescriptions and written advice/information on why an antibiotic was not prescribed (e.g. Non prescription pads). Studies show that this can be done within the same average consultation time while maintaining a high degree of patient satisfaction [12, 19-22].